INDIANA HEALTH COVERAGE PROGRAMS (IHCP) MISCELLANEOUS CARDIAC AGENTS PRIOR AUTHORIZATION REQUEST FORM



MDwise Fax to: (858) 790-7100 c/o MedImpact Healthcare Systems, Inc. Attn: Prior Authorization Department 10181 Scripps Gateway Court, San Diego, CA 92131 Phone: (808) 788-2949



	ay's Date / / / / / / / / / / / / / / / / / / /				uest will be returned**			
	ient's			Date of Birth				
Patient's Name				Prescriber's Name				
	Prescriber's IN License #				Specialty			
Pre	Prescriber's NPI #				Prescriber's Signature			
Re	turn Fax #			Return Phone	ne			
Ch	Check box if requesting retro-active PA			Date(s) of service requested for retro-active eligibility (if applicable):				
Note: Submit PA requests for retroactive claims (dates of service prior to eligibility determination, but within established eligibility timelines) with dates of service prior to 30 calendar days of submission separately from current PA requests (dates of service 30 calendar days or less and going forward).								
Requested Medication			Strength		Dosage Regimen			
	Requirements for Camzyos							
	Diagnosis of symptomatic obstructive hypertrophic cardiomyopathy (Provide documentation) ☐ Yes ☐ No							
	Left ventricular ejection fraction is greater than or equal to 55% (Provide documentation) ☐ Yes ☐ No							
	Left ventricular outflow tract (LVOT) gradient of 50 mm Hg or greater (Provide documentation) \square Yes \square No							
4. N	Member is 18 years of age or older ☐ Yes ☐ No							
5. N	Member is enrolled in Camzyos/mavacamten REMS program ☐ Yes ☐ No							
6. N								
C	channel blocker therapy ☐ Yes ☐ No							
	OR							
	Please provide medical rationale for the use of Camzyos (mavacamten) over beta-adrenergic blocker and non-dihydropyridine calcium channel blocker therapy							
7. F	Requested dose exceeds 15	 i mg/day □	Yes □ No					
	lote the following QL per strengt			g capsule – m	nax 1 capsule/day			

PA	Red	quirements for Corlanor (ivabradine) Tablet or Corlanor (ivabradine) Solution for Adults:			
	Sel	ect one of the following:			
	☐ Diagnosis of heart failure (Provide documentation)				
	$ullet$ Left ventricular ejection fraction is less than or equal to 35% (Provide documentation) \Box Yes \Box No				
		 Resting heart rate is greater than or equal to 70 beats per minute (Provide documentation) ☐ Yes ☐ No 			
		Diagnosis of inappropriate sinus tachycardia			
2.	Sel	lect one of the following: Member is currently maximized on beta-blocker dose			
		Drug/dose/date(s): Member has contraindication to beta-blocker use			
	Please explain:				
3.	. Select one of the following:				
	☐ Tablet Requested dose does not exceed 15 mg/day ☐ Yes ☐ No Note the following QL per strength: 5 mg, 7.5 mg, tablet – max 2 tablets/day				
		☐ Solution Requested dose does not exceed 15 mL/day ☐ Yes ☐ No			
		 Member is unable to swallow tablet formulation (Provide documentation) ☐ Yes ☐ No Note only approvable for a member who is 18 years of age or older and cannot swallow tablets 			
4.	Me	mber is 18 years of age or older □ Yes □ No			
PA	Red	quirements for Corlanor (ivabradine) Tablet or Corlanor (ivabradine) Solution for Pediatrics:			
		quirements for Corlanor (ivabradine) Tablet or Corlanor (ivabradine) Solution for Pediatrics: gnosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation)			
	Dia				
	Dia	ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation)			
1.	Dia	ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No			
1.	Dia Lef	Ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No t ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No			
 1. 2. 3. 4. 	Dia Lef Me	Ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No It ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No mber is in sinus rhythm (Provide documentation) □ Yes □ No			
 1. 2. 3. 4. 	Dia Lef Me	Ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No It ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No Important in sinus rhythm (Provide documentation) □ Yes □ No Sting heart rate is elevated (Provide documentation) □ Yes □ No Sect one of the following:			
 1. 2. 3. 4. 	Dia Lef Me	Ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No It ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No Important in sinus rhythm (Provide documentation) □ Yes □ No Institute of the following: Institute of the following: Institute of the following is described and institute of the following: Institute of the following: Institute of the following is described and institute of the following: Institute of the following is described and institute of the following: Institute of the following is described and institute of the f			
 1. 2. 3. 4. 	Diaa Leff Me Res	Ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No t ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No mber is in sinus rhythm (Provide documentation) □ Yes □ No sting heart rate is elevated (Provide documentation) □ Yes □ No lect one of the following: Member is 6 months through 17 years of age and ≥ 40 kg Request is for tablet formulation □ Yes □ No Requested dose does not exceed 15 mg/day □ Yes □ No Note the following QL per strength: 5 mg, 7.5 mg, tablet – max 2 tablets/day			
 1. 2. 3. 4. 	Diaa Leff Me Res	Ignosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No It ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No Important in sinus rhythm (Provide documentation) □ Yes □ No Institute selevated (Provide documentati			
 1. 2. 3. 4. 	Diaa Leff Me Res	Ingnosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No It ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No Important in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No			
 1. 2. 3. 4. 	Dia Lef Me Res Sel	Ingrosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No It ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No Important is in sinus rhythm (Provide documentation) □ Yes □ No Institute that is elevated (Provide documentation) □ Yes □ No Institute that is elevated (Provide documentation) □ Yes □ No Institute that is elevated (Provide documentation) □ Yes □ No Institute that is elevated (Provide documentation) □ Yes □ No Institute that is elevated (Provide documentation) □ Yes □ No Institute that is elevated (Provide documentation) □ Yes □ No Institute that is elevated (Provide documentation) □ Yes □ No Institute that is elevated documentation □ Yes □ No Institute that □ Yes □ No I			
 1. 2. 3. 4. 	Dia Lef Me Res Sel	Ingrosis of stable symptomatic heart failure due to dilated cardiomyopathy (Provide documentation) Yes □ No It ventricular ejection fraction is less than or equal to 45% (Provide documentation) □ Yes □ No Important material in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) □ Yes □ No Instituted in sinus rhythm (Provide documentation) Ins			

	 □ Member is 6 months through < 1 year of age and < 40 kg Requested dose does not exceed 0.2 mg/kg/dose twice daily
	☐ Yes ☐ No Weight:
PA	Requirements for Entresto (sacubitril-valsartan) sprinkle
1.	One of the following:
	☐ Member is less than 12 years of age and/or < 50 kg Weight:
	☐ Member is 12 years of age or older, ≥ 50 kg, and cannot swallow tablet formulation
2.	Prescriber attests to the following:
	 Member is/will NOT be using concomitant angiotensin converting enzyme (ACE) inhibitor or angiotensin II receptor blocker (ARB) therapy
PA	Requirements for Verquvo (vericiguat):
1.	Member is 18 years of age or older ☐ Yes ☐ No
2.	Diagnosis of chronic, symptomatic heart failure (Provide documentation) \square Yes \square No
3.	Left ventricular ejection fraction is less than or equal to 45% (Provide documentation) \square Yes \square No
4.	Select one of the following:
	☐ Member has been hospitalized for heart failure in the past 180 days (Provide documentation)
	☐ Member has received IV diuretics in the past 90 days (Provide documentation)
5.	For those of childbearing potential, documentation of a negative pregnancy test obtained within the past 60
	days is attached ☐ Yes ☐ No
6.	Requested dose exceeds 10 mg/day □ Yes □ No
	Note the following QL per strength: 2.5 mg, 5 mg, 10 mg tablet – max 1 tablet/day

CONFIDENTIAL INFORMATION

This facsimile transmission (and attachments) may contain protected health information from the Indiana Health Coverage Programs (IHCP), which is intended only for the use of the individual or entity named in this transmission sheet. Any unintended recipient is hereby notified that the information is privileged and confidential, and any use, disclosure, or reproduction of this information is prohibited.