

Smoking: Should I Take Medicine to Quit?

Making health decisions is part of life. These decisions can have a big effect on your health and happiness. Most people feel better about their health care when they take part in these decisions.

This information will help you understand your choices, whether you take part in the decision or ask your doctor to decide. Your decision should be based on the medical facts as well as your own feelings.


Your choices

- You can try to quit smoking without taking medicine.
- You can take medicine to help you quit. It can also help you manage cravings and withdrawal symptoms, such as anxiety or getting irritated easily.
- You can also combine some nicotine replacements, quit-smoking medicine, or other quit strategies such as counseling.

Key points in making your decision

Today, you have the choice of using medicines to improve your chances of quitting. You can use nicotine replacement products—like gum, patches, lozenges, nasal spray, and inhalers—that have nicotine in them. Or you can use medicines that do not have nicotine, such as varenicline (Chantix) or bupropion SR.

All of these medicines help ease your cravings and withdrawal symptoms. And they rarely cause serious side effects or addiction. These products do not have the unhealthy substances that are in cigarettes, such as tar and carbon



monoxide.

Think about the following when making your decision:

- Using medicines increases your chances of quitting smoking.
- Medicines and some nicotine replacement products need a prescription from your doctor.
- Some insurance companies may pay for all or part of the cost of quit-smoking medicines. If you do have to pay for it yourself, try to think about all the money you will save when you stop having to buy nicotine products.
- Nicotine replacement medicines like gum, lozenges, and inhalers can give quick relief for cravings.
- Nicotine gum, lozenges, and patches can be bought without a prescription.
- Nicotine patches do not give you quick relief for cravings. They are used once a day.

- The side effects of nicotine replacement medicines can include sore throat, upset stomach, skin rashes, and sleep problems.
- Dry mouth and sleep problems are the most common side effects of bupropion SR. Nausea and constipation are the most common side effects of varenicline.

Think about the facts and about what matters most to you. To help with your decision, take this information to your doctor to talk about your thoughts and feelings. Ask for more information if needed.

Reasons to take medicine to quit smoking	What to consider before choosing medicines
<ul style="list-style-type: none"> • You have cravings or withdrawal symptoms such as nervousness, feeling hungry, and feeling grumpy when you go too long without smoking. • You have tried to quit in the past without using medicine, but you started smoking again. • You want to make the best effort you can to quit. 	<ul style="list-style-type: none"> • If you have a history of seizures, mental health conditions, or heart disease, see your doctor before choosing which medicine to use. • If you may get pregnant, see your doctor before choosing which medicine to use.

Medicines have side effects and may not be right for everyone. Talk to your doctor about which medicine might be the best choice for you.

By using what you have just learned, you will be able to make a better decision about taking medicine to quit smoking.

- Learn the facts.
- Consider the benefits and risks of the different medicines.
- Ask about costs.
- Talk to your doctor.

Where can you find out more?

Smokefree.gov: 1-800-QUITNOW
(1-800-784-8669) www.smokefree.gov

American Cancer Society: 1-800-ACS-2345
(1-800-227-2345) www.cancer.org

National Cancer Institute:
1-800-4-CANCER (1-800-422-6237)
www.cancer.gov

American Lung Association: 1-800-LUNG-USA
(1-800-586-4872) www.lung.org