



Sickle Cell Disease: Splenic Sequestration

Table of Contents

- Overview
- Related Information
- Credits

Overview

Splenic sequestration is a problem with the spleen that can happen in people who have sickle cell disease.

It happens when a lot of sickled red blood cells get trapped in the spleen. The spleen can enlarge, get damaged, and not work as it should. When the spleen doesn't work well, a person is more likely to have serious, life-threatening infections with certain types of bacteria.

If splenic sequestration happens suddenly, it can be a life-threatening emergency.

Those affected

This condition is more common in infants and young children who have sickle cell disease. It may follow a respiratory infection.

In older children and adults, the spleen often doesn't work because of years of damage from sickled cells.

What are the symptoms?

Splenic sequestration causes sudden and severe anemia, with symptoms of sudden weakness, pale lips, rapid breathing, excessive thirst, belly pain, and rapid heartbeat. Splenic sequestration can happen in people who have sickle cell disease.

Related Information

- Sickle Cell Disease

Credits

Current as of: December 13, 2023

Author: Ignite Healthwise, LLC Staff (<https://www.healthwise.org/specialpages/legal/abouthw/en>)

Clinical Review Board (<https://www.healthwise.org/specialpages/legal/abouthw/en>)

All Healthwise education is reviewed by a team that includes physicians, nurses, advanced practitioners, registered dietitians, and other healthcare professionals.

Note: The "printer friendly" document will not contain all the information available in the online document. Some information (e.g. cross-references to other topics, definitions or medical illustrations) is only available in the online version.



This information does not replace the advice of a doctor. Ignite Healthwise, LLC, disclaims any warranty or liability for your use of this information.

© 2024 Ignite Healthwise, LLC.